

The Magnetospheres and Space Weather Environments of Extrasolar Planets


Optimized Strategies for Detecting Extrasolar Space Weather



Caltech



Gregg Hallinan

E-mail: gh@astro.caltech.edu



Sept 6th, 2017


Sunspot AR2673



2017/09/06 12:30


SDO/AIA 131 2017-09-06 00:12:56 UT

Severe storm conditions met at: 07/2350 UTC



G4

G4




- Flares – higher X-ray and ultraviolet radiation flux → photochemical reactions leading to significant atmospheric loss
- Coronal mass ejections (CMEs) – higher stellar wind flux → can erode atmosphere – eg. ion pick-up of a CO²-rich atmosphere



Magnetic activity can redefine habitability!

Stars out to 8 pc



Kirkpatrick et al. 2012




Small planets dominate planetary demographics and favor smaller stars (Howard et al. 2012)

Rocky planets are particularly frequent around M dwarfs (Dressing & Charbonneau 2013, 2015)

The nearest habitable planet likely orbits an M dwarf at 2.6 ± 0.4 pc

Trappist-1

Anglada-Escudé et al 2016



Credit: NASA/JPL-Caltech

Proxima b

Gillon et al 2016



Credit: ESO/M. Kornmesser





Image Credit: Chester Harman



West et al. 2008



Credit: KISS/Caltech

**Is magnetic activity important for defining habitability?
Can we directly detect CMEs and planetary magnetic fields?
Yes – with radio observations**

Low Frequency Radio Emission



**Type II radio emission
associated with CMEs**

**Planetary auroral
radio emission**


Strategy 1: Targeted Searches

Ongoing Searches for Stellar CMEs


Long bursts (>~1 hour)
Requires ongoing electron acceleration

Short bursts (sec - min)
Powered by individual flares?


AD Leo (M3.5)



UV Cet (M6)



YZ CMi (M4.5)



Time (min)


Villadsen, GH et al. 2018

- *Stellar dynamic spectroscopy a mature field (Bastian & Bookbinder 1987, Osten & Bastian 2006)*
- *Recent study – 21 bursts with ultra-wide bandwidth, no Type II bursts (Villadsen, GH et al. 2018)*


- *Need more sensitivity at lower frequencies!*

Exoplanet Searches

- Searches have been ongoing for > 30 years
- No detections
- See Lazio et al. 2009 for review



New Kids on the Block




MWA: 80-300 MHz



HERA: 50-250 MHz






LOFAR: 10-240 MHz



LWA: 10-90 MHz



Brown Dwarf Radio and Optical Aurorae



Hallinan et al. 2015, *Nature*, 523, 568

- Mostly detected with the VLA at GHz frequencies → kG magnetic fields
- see recent reviews by Pineda, GH and Kao 2017; Williams 2017

Radio Emission from a Candidate Free Floating Planet – SIMP0136



Kao, GH et al. 2016, 2018


- Brightest T dwarf in the northern hemisphere (Artigau et al. 2006)
- Carina Near moving group association - age of 200 Myr (Gagne et al. 2017)

Estimated mass of 12.7 ± 1.0 Jupiter masses – first radio exoplanet?

Magnetic fields ~ 3 kG

Gaia will find many more candidates for the VLA...

Targeted Searches from Space




L-ROLS - 100 lunar-based dipoles (Bob's talk) – scaled to Alpha Cen system


- Long-term monitoring of nearest candidate habitable exoplanet hosts (e.g. Alpha Cen system)
- **Can we detect solar-like CMEs on Alpha Cen AB and Proxima Centauri?**
- ***Do M dwarfs produce radio bursts (and CMEs) as energetic as the Sun?***
- ***Exoplanets detection via this method likely requires $>10^4$ dipoles***

Strategy 2: Multiplexed Searches

Space Weather Is highly Variable




Gallagher & D'Angelo 1981




Li, Zhang & Feng 2016

The OVRO-LWA: An Extrasolar Space Weather Telescope




Stokes I



Stokes V





○ GJ 725






○ 61 Cygni


○ Ross 248

○ Epsilon Eridani

○ UV Ceti

288-antenna performance: 500 mJy (10 s); 50 mJy (1 hr)
352-antenna performance: 100 mJy (10s); 5 mJy (1 hr); 1 mJy (1 day)





Monitors ~4,000 stellar/planetary systems out to 25pc


Anderson, GH et al. 2017

Multiplexed Searches from Space







Planetary radio emission subject to scaling laws for magnetic field strength and input solar wind power (e.g. see Farrell et al. 1999)

Proxima b






Adapted from
Burkhart & Loeb 2017

Contextual Data in the Search for Biosignatures



Strategy 3: Triggered Searches for Biosignatures

Triggered Alerts from a Lunar Array



Simulated high-resolution spectrum of Proxima Cen b with 0.1 TW auroral emission at 5577 Å (Luger et al. 2017)

Summary



Understanding the impact of stellar activity and the presence of planetary magnetic fields is becoming increasingly important for defining planetary habitability

Low frequency radio observations are key

The long-term future is from the lunar far-side

Targeted searches are computationally low-cost but limited

Multiplexed searches require significant in-situ computational resources

Triggered searches for biosignatures present an exciting possibility